

**NEUROLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Appropriate Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Stroke and Stroke Rehabilitation: Discharged on Antithrombotic Therapy	N/A	N/A	N/A	032	Percentage of patients aged 18 years and older with a diagnosis of ischemic stroke or transient ischemic attack (TIA) who were prescribed antithrombotic therapy at discharge	Effective Clinical Care	Process			American Academy of Neurology	X	-	-	-	-	X
Care Plan	N/A	N/A	0326	047	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan	Communication and Care Coordination	Process	X		National Committee for Quality Assurance	X	-	-	-	-	X
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	CMS69v5	N/A	0421	128	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter  Normal Parameters: Age 18 years and older BMI => 18.5 and < 25 kg/m2	Community/Population Health	Process			Centers for Medicare & Medicaid Services	X	-	X	X	-	X
Documentation of Current Medications in the Medical Record	CMS68v6	N/A	0419	130	Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Patient Safety	Process	X		Centers for Medicare & Medicaid Services	X	-	X	-	-	X
Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	CMS138v5	N/A	0028	226	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	Community/Population Health	Process			Physician Consortium for Performance Improvement	X	-	X	X	-	X
Epilepsy: Counseling for Women of Childbearing Potential with Epilepsy	N/A	N/A	1814	268	All female patients of childbearing potential (12 - 44 years old) diagnosed with epilepsy who were counseled or referred for counseling for how epilepsy and its treatment may affect contraception OR pregnancy at least once a year	Effective Clinical Care	Process			American Academy of Neurology	X	-	-	-	-	X
Dementia: Cognitive Assessment	CMS149v5	N/A	N/A	281	Percentage of patients, regardless of age, with a diagnosis of dementia for whom an assessment of cognition is performed and the results reviewed at least once within a 12 month period	Effective Clinical Care	Process			Physician Consortium for Performance Improvement	-	-	X	-	-	-

**NEUROLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Appropriate Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Dementia: Functional Status Assessment	N/A	N/A	N/A	282	Percentage of patients, regardless of age, with a diagnosis of dementia for whom an assessment of functional status is performed and the results reviewed at least once within a 12 month period	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Dementia: Neuropsychiatric Symptom Assessment	N/A	N/A	N/A	283	Percentage of patients, regardless of age, with a diagnosis of dementia and for whom an assessment of neuropsychiatric symptoms is performed and results reviewed at least once in a 12 month period	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Dementia: Management of Neuropsychiatric Symptoms	N/A	N/A	N/A	284	Percentage of patients, regardless of age, with a diagnosis of dementia who have one or more neuropsychiatric symptoms who received or were recommended to receive an intervention for neuropsychiatric symptoms within a 12 month period	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Dementia: Counseling Regarding Safety Concerns	N/A	N/A	N/A	286	Percentage of patients, regardless of age, with a diagnosis of dementia or their caregiver(s) who were counseled or referred for counseling regarding safety concerns within a 12 month period	Patient Safety	Process	X		American Academy of Neurology	-	-	-	-	-	X
Dementia: Caregiver Education and Support	N/A	N/A	N/A	288	Percentage of patients, regardless of age, with a diagnosis of dementia whose caregiver(s) were provided with education on dementia disease management and health behavior changes AND referred to additional resources for support within a 12 month period	Communication and Care Coordination	Process	X		American Academy of Neurology	-	-	-	-	-	X
Parkinson's Disease: Psychiatric Symptoms Assessment for Patients with Parkinson's Disease	N/A	N/A	N/A	290	All patients with a diagnosis of Parkinson's disease who were assessed for psychiatric symptoms (e.g., psychosis, depression, anxiety disorder, apathy, or impulse control disorder) in the last 12 months	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Parkinson's Disease: Cognitive Impairment or Dysfunction Assessment	N/A	N/A	N/A	291	All patients with a diagnosis of Parkinson's disease who were assessed for cognitive impairment or dysfunction in the last 12 months	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Parkinson's Disease: Rehabilitative Therapy Options	N/A	N/A	N/A	293	All patients with a diagnosis of Parkinson's Disease (or caregiver(s), as appropriate) who had rehabilitative therapy options (e.g., physical, occupational, or speech therapy) discussed in the last 12 months	Communication and Care Coordination	Process	X		American Academy of Neurology	-	-	-	-	-	X
Parkinson's Disease: Parkinson's Disease Medical and Surgical Treatment Options Reviewed	N/A	N/A	N/A	294	All patients with a diagnosis of Parkinson's disease (or caregiver(s), as appropriate) who had the Parkinson's disease treatment options (e.g., non-pharmacological treatment, pharmacological treatment, or surgical treatment) reviewed at least annually	Communication and Care Coordination	Process	X		American Academy of Neurology	-	-	-	-	-	X

**NEUROLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Appropriate Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	CMS22v5	N/A	N/A	317	Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated	Community/Population Health	Process			Centers for Medicare & Medicaid Services	X	-	X	-	-	X
Closing the Referral Loop: Receipt of Specialist Report	CMS50v5	N/A	N/A	374	Percentage of patients with referrals, regardless of age, for which the referring provider receives a report from the provider to whom the patient was referred	Communication and Care Coordination	Process	X		Centers for Medicare & Medicaid Services	-	-	X	-	-	-
Amyotrophic Lateral Sclerosis (ALS) Patient Care Preferences	N/A	N/A	N/A	386	Percentage of patients diagnosed with Amyotrophic Lateral Sclerosis (ALS) who were offered assistance in planning for end of life issues (e.g., advance directives, invasive ventilation, hospice) at least once annually	Person and Caregiver-Centered Experience and Outcomes	Process	X		American Academy of Neurology	-	-	-	-	-	X
Tobacco Use and Help with Quitting Among Adolescents	N/A	N/A	N/A	402	The percentage of adolescents 12 to 20 years of age with a primary care visit during the measurement year for whom tobacco use status was documented and received help with quitting if identified as a tobacco user	Community/Population Health	Process			National Committee for Quality Assurance	-	-	-	-	-	X
Opioid Therapy Follow-up Evaluation	N/A	N/A	N/A	408	All patients 18 and older prescribed opiates for longer than six weeks duration who had a follow-up evaluation conducted at least every three months during Opioid Therapy documented in the medical record	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Documentation of Signed Opioid Treatment Agreement	N/A	N/A	N/A	412	All patients 18 and older prescribed opiates for longer than six weeks duration who signed an opioid treatment agreement at least once during Opioid Therapy documented in the medical record.	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Evaluation or Interview for Risk of Opioid Misuse	N/A	N/A	N/A	414	All patients 18 and older prescribed opiates for longer than six weeks duration evaluated for risk of opioid misuse using a brief validated instrument (e.g. Opioid Risk Tool, SOAPP-R) or patient interview documented at least once during Opioid Therapy in the medical record	Effective Clinical Care	Process			American Academy of Neurology	-	-	-	-	-	X
Overuse Of Neuroimaging For Patients With Primary Headache And A Normal Neurological Examination	N/A	N/A	N/A	419	Percentage of patients with a diagnosis of primary headache disorder whom advanced brain imaging was not ordered	Efficiency and Cost Reduction	Efficiency	X	X	American Academy of Neurology	X	-	-	-	-	X
Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	N/A	N/A	2152	431	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 24 months AND who received brief counseling if identified as an unhealthy alcohol user	Community/Population Health	Process			Physician Consortium for Performance Improvement	-	-	-	-	-	X

**NEUROLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Appropriate Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Quality of Life Assessment For Patients With Primary Headache Disorders	N/A	N/A	N/A	435	Percentage of patients with a diagnosis of primary headache disorder whose health related quality of life (HRQoL) was assessed with a tool(s) during at least two visits during the 12 month measurement period AND whose health related quality of life score stayed the same or improved	Effective Clinical Care	Outcome	X		American Academy of Neurology	X	-	-	-	-	X