

**OBSTETRICS & GYNECOLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Approp Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Care Plan	N/A	N/A	0326	047	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan	Communication and Care Coordination	Process	X		National Committee for Quality Assurance	X	-	-	-	-	X
Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older	N/A	N/A	N/A	048	Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months	Effective Clinical Care	Process			National Committee for Quality Assurance	X	-	-	-	-	X
Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older	N/A	N/A	N/A	050	Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months	Person and Caregiver-Centered Experience and Outcomes	Process	X		National Committee for Quality Assurance	X	-	-	-	-	X
Preventive Care and Screening: Influenza Immunization	CMS147v6	N/A	0041	110	Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization	Community/Population Health	Process			Physician Consortium for Performance Improvement	X	-	X	X	-	X
Breast Cancer Screening	CMS125v5	N/A	2372	112	Percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.	Effective Clinical Care	Process			National Committee for Quality Assurance	X	-	X	X	-	X
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	CMS69v5	N/A	0421	128	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter  Normal Parameters: Age 18 years and older BMI => 18.5 and < 25 kg/m2	Community/Population Health	Process			Centers for Medicare & Medicaid Services	X	-	X	X	-	X

**OBSTETRICS & GYNECOLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Approp Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Documentation of Current Medications in the Medical Record	CMS68v6	N/A	0419	130	Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Patient Safety	Process	X		Centers for Medicare & Medicaid Services	X	-	X	-	-	X
Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	CMS138v5	N/A	0028	226	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	Community/Population Health	Process			Physician Consortium for Performance Improvement	X	-	X	X	-	X
Controlling High Blood Pressure	CMS165v5	N/A	0018	236	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90mmHg) during the measurement period	Effective Clinical Care	Intermediate Outcome	X		National Committee for Quality Assurance	X	-	X	X	-	X
Biopsy Follow-Up	N/A	N/A	N/A	265	Percentage of new patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician	Communication and Care Coordination	Process	X		American Academy of Dermatology	-	-	-	-	-	X
Cervical Cancer Screening	CMS124v5	N/A	0032	309	Percentage of women 21-64 years of age who were screened for cervical cancer using either of the following criteria: * Women age 21-64 who had cervical cytology performed every 3 years * Women age 30-64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years	Effective Clinical Care	Process			National Committee for Quality Assurance	-	-	X	-	-	-
Chlamydia Screening for Women	CMS153v5	N/A	0033	310	Percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement period	Community/Population Health	Process			National Committee for Quality Assurance	-	-	X	-	-	-
Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	CMS22v5	N/A	N/A	317	Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated	Community/Population Health	Process			Centers for Medicare & Medicaid Services	X	-	X	-	-	X

**OBSTETRICS & GYNECOLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Approp Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Closing the Referral Loop: Receipt of Specialist Report	CMS50v5	N/A	N/A	374	Percentage of patients with referrals, regardless of age, for which the referring provider receives a report from the provider to whom the patient was referred	Communication and Care Coordination	Process	X		Centers for Medicare & Medicaid Services	-	-	X	-	-	-
Tobacco Use and Help with Quitting Among Adolescents	N/A	N/A	N/A	402	The percentage of adolescents 12 to 20 years of age with a primary care visit during the measurement year for whom tobacco use status was documented and received help with quitting if identified as a tobacco user	Community/Population Health	Process			National Committee for Quality Assurance	-	-	-	-	-	X
Osteoporosis Management in Women Who Had a Fracture	N/A	N/A	0053	418	The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis in the six months after the fracture	Effective Clinical Care	Process			National Committee for Quality Assurance	X	-	-	-	-	X
Performing Cystoscopy at the Time of Hysterectomy for Pelvic Organ Prolapse to Detect Lower Urinary Tract Injury	N/A	N/A	2063	422	Percentage of patients who undergo cystoscopy to evaluate for lower urinary tract injury at the time of hysterectomy for pelvic organ prolapse	Patient Safety	Process	X		American Urogynecologic Society	X	-	-	-	-	X
Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	N/A	N/A	2152	431	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 24 months AND who received brief counseling if identified as an unhealthy alcohol user	Community/Population Health	Process			Physician Consortium for Performance Improvement	-	-	-	-	-	X
Proportion of Patients Sustaining a Bladder Injury at the Time of any Pelvic Organ Prolapse Repair	N/A	N/A	N/A	432	Percentage of patients undergoing any surgery to repair pelvic organ prolapse who sustains an injury to the bladder recognized either during or within 1 month after surgery	Patient Safety	Outcome	X		American Urogynecologic Society	-	-	-	-	-	X
Proportion of Patients Sustaining a Bowel Injury at the time of any Pelvic Organ Prolapse Repair	N/A	N/A	N/A	433	Percentage of patients undergoing surgical repair of pelvic organ prolapse that is complicated by a bowel injury at the time of index surgery that is recognized intraoperatively or within 1 month after surgery	Patient Safety	Outcome	X		American Urogynecologic Society	-	-	-	-	-	X
Proportion of Patients Sustaining a Ureter Injury at the Time of any Pelvic Organ Prolapse Repair	N/A	N/A	N/A	434	Percentage of patients undergoing pelvic organ prolapse repairs who sustain an injury to the ureter recognized either during or within 1 month after surgery	Patient Safety	Outcome	X		American Urogynecologic Society	-	-	-	-	-	X

**OBSTETRICS & GYNECOLOGY 2017 - 18**

Measure Title	Measure Number				Measure Description	NQS Domain	Measure Type	High Priority	Approp Use	Primary Measure Steward	Data Submission Method(s)					
	eMeasure ID	eMeasure NQF	NQF	Quality No. (Q#)							Claims	CSV	EHR	CMS Web Interface	Admin Claims	Registry
Non-Recommended Cervical Cancer Screening in Adolescent Females	N/A	N/A	N/A	443	The percentage of adolescent females 16–20 years of age who were screened unnecessarily for cervical cancer	Patient Safety	Process	X	X	National Committee for Quality Assurance	-	-	-	-	-	X
Chlamydia Screening and Follow Up	N/A	N/A	N/A	447	The percentage of female adolescents 16 years of age who had a chlamydia screening test with proper follow-up during the measurement period	Community/Population Health	Process			National Committee for Quality Assurance	-	-	-	-	-	X
Appropriate Workup Prior to Endometrial Ablation	N/A	N/A	0567	448	Percentage of women, aged 18 years and older, who undergo endometrial sampling or hysteroscopy with biopsy before undergoing an endometrial ablation	Patient Safety	Process	X		Centers for Medicare & Medicaid Services	-	-	-	-	-	X